

D 31617

(Pages : 2)

Name

Reg. No.

THIRD SEMESTER M.Sc. DEGREE EXAMINATION, DECEMBER 2012

(Non-CUCSS)

Mathematics

Paper XII—FUNCTIONAL ANALYSIS—I

(2002 Admissions)

Maximum : 80 Marks

Time : Three Hours

Part A

*Answer all questions.
Each question carries 4 marks.*

- I. (a) Define Cauchy sequence and show that every Cauchy sequence is bounded. Is the converse true ? Justify your answer.
- (b) Prove that among all the normed spaces $L^p([0, 1])$; $1 \leq p < \infty$, only the space $L^2([0, 1])$ is an inner product space.
- (c) Show that the linear space C_0 cannot be a Banach space in any norm.
- (d) Let $\|\cdot\|$ be a complete norm on $C([a, b])$ such that if $\|x_n - x\| \rightarrow 0$, then $x_n(t) \rightarrow x(t)$ for every $t \in [a, b]$. Show that $\|\cdot\|$ is equivalent to the sup norm on $C([a, b])$.
- (4 x 4 = 16 marks)

Part B

*Answer any four questions without omitting any unit.
Each question carries 16 marks.*

UNIT I

- II. (a) Show that the intersection of a finite number of dense open subsets of a metric space X is dense in X .
- (b) Let T be a compact metric space and $E \subset C(T)$. Suppose that E is bounded and equicontinuous at each $t \in T$. Show that E is totally bounded in the sup metric on $C(T)$.
- III. (a) Show that for $1 < p < \infty$, the metric space $L^p(E)$ is complete for any measurable subset E of \mathbb{R} .
- (b) Let x be a continuous k -valued function on $[-\pi, \pi]$ such that $x(n) = x(-\pi)$. Show that the sequence of arithmetic means of the partial sums of the Fourier series of x converges to x uniformly on $[-\pi, \pi]$.

Turn over

IV. (a) State and prove Riesz Lemma.

(b) Let X be a normed space. Show that X is finite dimensional if every closed and hour 1 subset of X is compact.

UNIT II

V. (a) Show that every linear map on a finite dimensional normed space is continuous.

(b) Show that a linear functional f in a normed space X is continuous iff the zero space $Z(f)$ is closed in X .

(c) Let X be a normed space and $P \in BL(X)$ satisfy $P^2 = P$. Show that $\|P\| = 0$ or $\|P\| = 1$.

VI. (a) Let X be a normed space over k , Y be a subspace of X and $g \in Y'$. Show that there is some

$$f \in X' \text{ such that } f|_Y = g \text{ and } \|f\| = \|g\|.$$

(b) Show that there exists a linear functional f on V such that

$$\|f\|^{-1} = f(a) \text{ and } f(x) = f(x) \text{ for all } x \in V \text{ where } a = \dots \text{ and } T(x)(j) = x(j+1) \text{ for } j=1, 2, \dots$$

VII. (a) Let $\{u_a\}$ be an orthonormal set in a Hilbert space H . Show that $\{u_a\}$ is an orthonormal basis for H iff $x \in H$ and $(x, u_a) = 0$ for all a , then $x = 0$.

(b) Let $H = L^2([0,1])$. Show that $\{1, \sqrt{2} \cos \pi t, \sqrt{2} \cos 2\pi t, \dots\}$ is an orthonormal basis for H .

UNIT III

VIII. (a) Show that a normed space X is a Banach space iff every absolutely summable series of elements in X is summable in X .

(b) Show that every normed can be embedded as a dense subspace of a Banach space.

IX. (a) Let X be a normed space and E be a subset of X . Show that E is bounded in X if $f(E)$ is bounded in \mathbb{R} for every $f \in X'$.

(b) Let $1 \leq p < \infty$ and $X = C_0$ with the norm $\| \cdot \|_p$. For $n = 1, 2, \dots$, let $f_n(x) = nx(n)$; $x \in X$.

Show that $f_n(x) \rightarrow 0$ for every $x \in X$, but $\|f_n\| \rightarrow \infty$.

X. (a) State and prove closed graph theorem.

(b) Show that the closed graph theorem may not hold for normed spaces.

(4 x 16 = 64 marks)