C 60050

(Pages: 3)

Name.....

Reg. No.....

SIXTH SEMESTER B.Sc. DEGREE EXAMINATION, MARCH 2019

(CUCBCSS)

Mathematics

MAT 6B 11-NUMERICAL METHODS

Time : Three Hours

Maximum : 120 Marks

Section A

Answer all the **twelve** questions. Each question carries 1 mark.

1. Give an example of a transcendental function.

2. What do you mean by complete pivoting ?

3. What is the advantage of Gauss Jordan method over Gauss Elimination method ?

4. Write Newton's forward difference interpolation formula.

5. Write Lagrange's interpolation formula of degree n.

- 6. State Trapezoidal rule.
- 7. What is meant by ill-conditioned system of equations ?
- 8. What is a differential equation ?
- 9. Find the degree and order of differential equation $y^1 + ay^2 = 0$.

10. What is interpolation?

11. What is homogeneous equation ?

12. Give an example of linear function.

$(12 \times 1 = 12 \text{ marks})$

Section B

Answer any **ten** out of fourteen questions. Each question carries 4 marks.

- 13. Use the method of fixed point iteration to find a positive root, between 0 and 1, of the equation $xe^{x} = 1$.
- 14. Find a real root of the equation $f(x) = x^3 x 1 = 0$.
- 15. Evaluate $e^{1.24}$, given that $e^{1.1} = 3.0042$ and $e^{1.4} = 4.0552$.
- 16. State Simpson's 1/3rd rule.

Turn over

17. Explain the limitations of using Newton-Raphson's method.

18. Construct a divided difference table for 4 data points.

19. State the formula of Picard's method to solve the differential equation of type dy/dx = f(xy).

20. Comment the accuracy of Euler's method.

21. What is triangularisation of equations?

22. State the second order Newton's divided difference interpolation polynomial.

23. What are the limitations of Taylor's series method.

24. Write the fourth order Runge-Kutta formula.

25. By the matrix inversion method solve :

$$2x + y = 1$$
$$x + 3y = 2.$$

26. What is the difference between Gauss elimination and Gauss-Jordan method ?

 $(10 \times 4 = 40 \text{ marks})$

Section C

Answer any **six** out of nine questions. Each question carries 7 marks.

27. Solve the system by using Gauss-Jordan method.

$$2x_1 + 4x_2 - 6x_3 = -8$$

$$x_1 + 3x_2 + x_3 = 10$$

$$2x_1 - 4x_2 - 2x_3 = -12.$$

28. Using linear interpolation formula estimate the square root of 2.5 :

X : 1 2 3 4 5 \sqrt{x} : 1 1.4142 1.7321 2 2.2361

29. The table below gives the values of distance travelled by a car at various time intervals during the initial running :

Time (s)	:	5	6	7	8	9
Distance (km)	:	10	14.5	19.5	25.5	32

Estimate velocity at time t = 5, t - 7 and t = 9.

30. Evaluate the $\int_{a}^{b} x^{3} + 1$ by using Trapezoidal rule for the (1, 2) and (1, 1.5).

31. Use Taylor's method to solve the equation y' = x² + y² for x = 0.25 and x = 0.5 given y (0) = 1.
32. Obtain a polynomial using Lagrange formula :

x : 0 1 2 3 $e^{x}-1$: 0 1.7183 6.3891 19.0855

Use the polynomial to estimate the value of $e^{1.5}$.

- 33. Find the root of the equation $x^2 3x + 2$ in the vicinity of x = 0 using Newton's Raphson method.
- 34. Explain the principle of false position method.
- 35. Given the equation $dy/dx = 3x^2 + 1$ with y(1) = 2. Find y(2) by Euler's method using h = 0.5 and h = 0.25.

 $(6 \times 7 = 42 \text{ marks})$

Section D

Answer any **two** out of three questions. Each question carries 13 marks.

36. (a) Use bisection method to find the root of the equation $x^2 - 4x - 10 = 0$.

- (b) Use false position method to determine the root of the equation $f(x) = x^2 x 2 = 0$ in the range 1 < x < 3.
- 37. (a) Solve the system by using Gauss-Seidel method :

$$2x_1 + x_2 + x_3 = 5$$

$$3x_1 + 5x_2 + 2x_3 = 15$$

$$2x_1 + x_2 + 4x_3 = 8.$$

(b) Solve the system by using Triangular factorization method :

$$3x_1 + 2x_2 + x_3 = 10$$

$$2x_1 + 3x_2 + 2x_3 = 14$$

$$x_1 + 2x_2 + 3x_3 = 14.$$

38. Use Runge-Kutta method to estimate y(0.4) when $y'(x) = x^2 + y^2$ with y(0) = 0 and assume h = 0.2.

 $(2 \times 13 = 26 \text{ marks})$